

Check for
Updates

EVALUATION OF INDUSTRIAL FRUITS WASTE CATALYTIC REACTIONS POTENTIAL WITH IONIC LIQUIDS

¹*Sanjay Kumar Pandey, ²Dr. Pankaj Gupta

*Correspondence: Sanjay Kumar Pandey

***Detailed author information and related declarations are provided in the final section of this article.*

Article Publication Details

This article is published in the **International Journal of Multidisciplinary Research and Bulletin**, ISSN 3108-1428 (Online) Volume 5 Issue 1 (Jan – Feb) 2026.

ABSTRACT

Fruit forests, veggies, decorative plants, root tubers, medicinal normal natural natural herbs, fragrant flowers, normal natural herbs, and plantation crops could all be grown in India's diverse agro environment. After usage, good fresh fruit peels certainly are an annoyance to the environment as solid waste. The top, actual, and chemical characteristics of commonly offered volume this is certainly fruit peels (FP) (such for instance banana, tangerine, citrus, lemon, and jackfruit) are examined in this analysis to make a detailed suggestion because of their monetization. After practices were used to characterize each FP: proximate and evaluation that is porosity that is particle that is ultimate, amount thickness, point of zero fees (pHpzc), area pH, surface expenses, liquid consumption capability, BET surface, checking electron microscopy, Fourier transforms infrared spectroscopy and TGA/thermogravimetric derivatives. The BET area of FP is minimal, varying between 0.60 and 1.2m²/g. The pH and top which are pHpzc of orange-peel (OP), citrus peel (CP), lemon peel (LP), and jackfruit skins (JFP) are in the 3–4 range. The value that is pHpzc nearer to 7 compared to the certain area that's sure of a banana peel (BP). Your order of area acidity is really as employs: OP>LP>CP>JFP>BP. The security of FPs below 150°C is seen from TG curves. The result could be really healing for logical design whenever FP is employed as being a substrate for bioactive substances, phenolic anti-oxidants, organic acids, enzymes, biofertilizers, energy production, and adsorbents.

Keywords: Green chemistry, Ionic compounds, Fruit peel waste, Catalyst, etc.

Introduction

The increased loss of fruit and waste isn't only a waste of foodstuffs, but additionally a waste of essential sources of land this is certainly including water, fertilizers, chemical compounds. As these huge volumes of wasted food, things disintegrate in landfills and release greenhouse this is certainly harmful (Venkat 2011; Valerio yet others 2017), they contribute to huge ecological problems. Fruit and veggie handling products, which are usually followed by domestic rubbish, produce the most trash into the environment. Fresh fruits perform a component that is essential to our nourishment as well as in-person life, and for that reason of this expanding international population and changing diet preferences, demand for such crucial meals products has increased dramatically (Schieber and others 2001; Valerio as well as others 2017). Citrus accounts for 124.73 million tons which can be metric MMT), bananas for 114.08 MMT, apples for 84.63 MMT, grapes for 74.49 MMT, mangoes, mango steins, and guavas for 45.22 MMT, and pineapples for 25.43 MMT of global fresh fruit production.

REVIEW OF LITERATURE:

“Plant secondary metabolites labeled as phenolic compounds are responsible for sensory features and subscribe to the high quality this is certainly health of and vegetables, among other things (Tomás-Barberán as well as others 2000; Lapornik among others 2005). Phenolic compounds are probably the most diverse categories of bioactive chemicals, having a range this is certainly wide of results (Popa among others 2008; Ignat and others 2011). Obtained 1 or more fragrant rings within their fundamental construction, along with 1 or higher hydroxyl teams (Balasundram and colleagues 2006), and some program action that is antioxidant Heim and others 2002). Amongst others, tend to be polyphenolic chemical substances categorized into many courses. The citrus company makes an important wide range of seeds and peel waste, which accounts fully for nearly 50% of the good fresh fruit this is certainly total Bocco yet others 1998). Citrus garbage is a supply this is certainly abundant with compounds because citrus peel includes a greater concentration of polyphenols compared to the section that is edible of fruit (Balasundram while others 2006).

The peels of numerous other fruits have now been found to obtain better items that tend to be phenolic to your parts which are delicious alongside citrus. Banana pulp (Musa cavendish) had been discovered to have 232 mg/100 DM that is g of substances, that is just around 25% for the amount found in the peel (Someya among others 2002). As well as chemical compounds that are phenolic banana skins had greater quantities of catecholamines, dopamine, and 1-dopa (González-

Montelongo among others 2010). The skins of pomegranates have 249.4 Mg/g chemicals being phenolic whereas the pulp has 24.4 mg/g (Li among others 2006). The skins of eight Clingstone peach varieties had been discovered to possess a lot more than 2.0 to 2.5 times the amount of phenolic chemicals found in the skin this is certainly edible change and others 2000). Because phenolic mixture extraction is based on the technique used, ultrasound-assisted extraction can enhance phenolic compound recovered from (*Citrus reticulata* L.) peel waste by four times when compared with maceration (Safdar yet others 2016)”[1].

MATERIALS AND METHODS

Material

Wastes from a local market in the District include banana fruit peels, orange fruit peels, citrus fruit peels, lemon fruit peels, and jackfruit peels. The peeling wastes are first washed (at least 5 times) to eliminate any dirt or moisture that has adhered to them. At room temperature (25°C), they are stored and used for further research.

Sample preparation

Skins from bananas, oranges, citrus fruits, lemons, and jackfruit were purchased at a fresh fruit that is regional in India's district degree. From then on, the leaves, sticks, along other things are free removed. The peels were oven-dried at $70\pm 2^{\circ}\text{C}$ till they reached a frequent fat after precisely rinsing with plain tap water to eliminate any real imperfections at first glance. It had been sieved and powdered at the last end of this process.

RESULTS AND DISCUSSION:

Physicochemical characterization

More or less and ultimately researches of each peel had been done to find out properties such for instance porosity, the thickness of particles, the thickness of bulk, zero charge point, area ph, surface charges, liquid absorption capability, and wayer surface. In inclusion, samples were scanned with electron microscope (SEM), Fourier infra round transforms (FTIR which is certainly spectroscopy and TG/DTG methods. This study's characterization methodology and instrumentation techniques are described in depth.

Table 1. Proximate and ultimate analysis of Fruits peel

Proximate analysis of fruits peel					Ultimate analysis of fruits peel				
FP	Moisture (%)	Ash (%)	Volatile matter (%)	Fixed carbon (%)	N (%)	C (%)	H (%)	S (%)	O % (by diff.)
BP	9.68	5.11	84.38	0.83	1.33	40.22	6.14	0.1	52.22
OP	7.84	5.19	85.9	1.07	1.16	38.87	6.19	0.11	53.64
CP	7.52	4.38	85.68	2.42	0.68	38.56	6.2	0.1	54.55
LP	6.23	5.46	86.18	2.13	1.24	40.28	5.96	0.19	52.25
JFP	6.54	6.23	86.22	1.01	0.96	40.11	5.86	0.12	53.08

Physical properties of FP

Physical properties such as porosity, particle density, bulk water-absorption, and density capability of FP was in fact determined (Table 2). As summarized in dining table 2, CP contains the amount density that is most affordable however the water-absorption capability that is highest in comparison to various other FPs. OP contains the bulk depth that is greatest, nevertheless the most percentage that is affordable of. The differences in bulk densities for the FPs are usually due mainly to variants in particle dimensions, particle shape, or both. The volume regarding the atmosphere this is certainly entrapped by having an rise in porosity.

Table 2. Physical properties of various Fruits peel.


Fruit Peel	Water absorption capacity (ml/g)	Particle density (g/cc)	Bulk density (g/cc)	Porosity (%)
Jack Fruit Peel	5.2	0.82	0.41	42.36
Banana Peel	5.1	0.89	0.39	56.41
Lemon Peel	5.9	0.98	0.49	48.97
Orange Peel	5.4	0.89	0.53	41.13
Citrus Peel	6.7	1.06	0.38	63.91

BET surface area of fruit peel

The top FP is porous and rough as a whole, as reduction through the SEM pictures shown and discussed into the specific area that is after. The BET area of BP (0.65 m²/g), OP (1.03 m²/g), CP (1.14 m²/g), LP (0.70 m²/g) and JFP (1.15 m²/g) is really paid off, in comparison to services and products being siliceous. The BET area of BP is within the number of 1.2–13 m²/g, whereas when compared with OP is within the number of 2.14– 47.03. The liner that is external is low is possible because of the functional complexity of degassing lignocellulosic samples. Even moreover, the outer lining that is low is just a characteristic function of carbonaceous products.

SEM of Fruit peel

Figure 1 Scanning Electron Micrographs of (a) BP, (b) OP, (c) CP, (d) LP, and (e) JFP.



Utilisation of Fruit Peels as catalyst in Ionic Reaction

Observance of this is certainly microscopic particle geometry and architectural variants associated with the area. The surface of FP is uneven, harsh, and possesses some materials; also, the particle form and sizes can be various. The particles vary in form from spherical to flakes being very long. Some macropores and micropores are observed on the top besides. The top BP is rough and permeable (Figure 1 a). Even though the particular section of OP (Figure 1 b) can also be harsh, it

also contains some materials being spread through the entire area. The real wide range of follicles regarding the OP surface is less when compared with BP. CP (Figure 1 c) reveals a location that is irregularly minimal little pores. The top that is LP Figure 1 d) is normally asymmetrical and highly porous. In addition, its area (LP) possesses larger quantity of skin pores in comparison with other FPs. JFP features very long flakes like construction (Figure 1 e) containing a place that is harsh pores being little.

Fruit Peels are separated and discarded in municipal landfills after the edible portion has been consumed. This results in significant pollution as well as disposal (solid waste management) issues. As a dual purpose: (i) creating money from waste, and (ii) reducing solid waste.

Imines, often known as Schiff basics, are compounds with an azomethine group ($\text{HC}=\text{N}$) with the formula that is basic. For the synthesis of Schiff bases, certain eco-friendly and cost-effective normal catalysts such as grape juice (*Vitis limetta*), nice lemon liquid, and aqueous herb of mango (*Mangifera indica*) are utilized.

Ionic Liquids

Ionic liquids (ILs) are molten salts of anionic character that contain both cationic and anionic species and have a melting point below 100°C . Walden created the first ionic liquid $[\text{EtNH}_3][\text{NO}_3]$ (m.p. 12°C) in 1914, which began the history of ILs. Hurley and Wier created low melting salts containing chloroaluminate ions in 1951 for low-temperature aluminum electroplating. Due to their moisture sensitivity, these chloroaluminate salts did not garner much interest for applications. Wilkes group took another step forward in 1992 by synthesizing environment and water steady 1-ethyl-3-methylimidazolium-based fluids which can be ionic various anions. Davis and colleagues developed a novel class of ionic liquids dubbed "**functionalized ionic liquids (FILS)**" based on cations generated from the antifungal medication miconazole in 1998.

Schiff bases analysis

Preparation of catalyst:

Locally sourced banana fruit skins, orange good fresh fruit peels, citrus fruit peels, lemon-fresh fruit skins, and jackfruit peels are broken into fruit juice and blocked with cotton to acquire fluid juice from all good fresh fruit skins. While sweet lime fresh fruits are peeled using a knife plus the peels are squeezed to the liquid to secure a size that is semisolid that is then purified with cotton to have liquid to be used as being a catalyst.


Catalyst preparation from Fruit Peel

A fresh fruit that is neighborhood supplied the banana skins. The banana peel was dried in a range of 373K for 12 hours and paid down to dust using an agate mortar after numerous rinses with distilled liquid to get rid of solid impurities. The catalysts were created by calcining powder at various temperatures (500–800°C) in a muffle furnace for 2 hours at a heating rate of 2°C per min. All of the fruit peels are preparing themselves as if they were banana fruit peel catalysts. A clear fluid component is obtained through the upper shell of an unripened fruit and the fruit peel (5 gm) is cooked with 100 ml of water and then refreshed and filtered by the muslin fabric.

Synthesis of Schiff bases

Synthesis of Schiff bases using Banana fruit peel, Orange fruit peel, citrus fruit peel, Lemon fruit peel, and Jackfruit peel juice as catalysts and an aqueous extract of unripe mango as an aqueous extract in a solvent-free environment:

In several beakers, an equimolar amount of benzaldehyde was mixed with an equal amount of aniline. Natural acid catalysts, such as banana fruit peel, orange fruit peel, citrus fruit peel, lemon fruit peel, and jackfruit peel juice, “are added in several amounts (0.5 ml, 1 ml, 1.5 ml, 2.0 ml, 2.5 ml) to those response mixtures then held for 5-10 minutes. Each reaction combination had been stirred for another 2 to 4 mins at room temperature. After the response was completed, a pale yellow crude that is solid appeared, which had been washed with distilled liquid and processed by recrystallization with ethanol”[2].



The same technique is used with every catalyst and good fresh fruit peel herb that is aqueous. The open capillary method was used to determine the melting point of the products, which was then identified and purified using TLC and validated using mass spectra. We could observe that a nucleophilic attack of the primary amine in the carbonyl carbon produces a hydroxyl molecule, that is then dehydrated to create Schiff basics. If we consider just how aldehydes and amines come to be Schiff bases” [2]. The pace at which water is removed from the reaction mixture affects the production of Schiff bases in the second stage.

CONCLUSION:

The study had been performed to experience the characterization this is certainly physicochemical of peel waste, banana peel waste, citrus peel waste, lemon peel waste, and jackfruit peel waste. Various catalyst preparation by using various fruits peels waste for green synthesis of organic compounds (Schiff base, Knoevenagel condensation, multicomponent reaction) from banana peels, orange peels, citrus peels, lemon peels, and jackfruit peel wastes. In this research, we also studied the characterization of ionic liquids in catalytic reactions.

Article History

Received: 25-Jan-2026

Accepted: 01-Feb-2026

Published: 03-Feb-2026

Article Publication Details (rpt*)

This article is published in the [International Journal of Multidisciplinary Research and Bulletin](#), ISSN 3108-1428 (Online). In Volume 5 Issue 1 (Jan – Feb) 2026

The journal is published and managed by [IRPG](#).

Copyright © 2026, Authors retain copyright. Licensed under the Creative Commons Attribution 4.0 International License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. <https://creativecommons.org/licenses/by/4.0/> (CC BY 4.0 deed)

Acknowledgements

We sincerely thank the editors and the reviewers for their valuable suggestions on this paper.

Funding

The authors declare that no funding was received for this work.

Data availability

No datasets were generated or analyzed during the current study.

Declarations**Ethics approval and consent to participate**

The author(s) declare that it is not applicable.

Consent for publication

The author(s) declare that this is not applicable.

Competing interests

The author(s) declare that they have no competing interests.

Author details

Sanjay Kumar Pandey^{1*}

^{1*}Research Scholar, Department of Chemistry, Sunrise University, Alwar, Rajasthan, India.

Dr. Pankaj Gupta²

²Faculty of Science, Sunrise University, Alwar, Rajasthan, India.

References

1. Galanakis CM. 2012. Recovery of high added-value components from food wastes: conventional, emerging technologies and commercialized applications. *Trends Food Sci Technol* 26: 68– 87.
2. Rudra SG, Nishad J, Jakhar N, Kaur C. 2015. Food industry waste: mine of nutraceuticals. *Intl J Sci Environ Technol* 4: 205– 29.
3. Soong YY, Barlow PJ. 2004. Antioxidant activity and phenolic content of selected fruit seeds. *Food Chem* 88: 411– 7.
4. Gorinstein S, Martín-Belloso O, Park YS, Haruenkit R, Lojek A, Číž M, Caspi A, Libman I, Trakhtenberg S. 2001a. Comparison of some biochemical characteristics of different citrus fruits. *Food Chem* 74: 309– 15.
5. Gorinstein S, Zachwieja Z, Folta M, Barton H, Piotrowicz J, Zemser M, Martín-Belloso O. 2001b. Comparative contents of dietary fiber, total phenolics, and minerals in persimmons and apples. *J Agric Food Chem* 49: 952– 7.
6. Lapornik B, Prošek M, Wondra AG. 2005. Comparison of extracts prepared from plant by-products using different solvents and extraction time. *J Food Engr* 71: 214– 22.
7. Tomás-Barberán FA, Ferreres F, Gil MI. 2000. Antioxidant phenolic metabolites from fruit and vegetables and changes during postharvest storage and processing. In: A-U Rahman, editor. *Studies in natural products chemistry*. Vol. 23. Amsterdam, The Netherlands: Elsevier Science. p 739– 95.
8. Popa VI, Dumitru M, Volf I, Anghel N. 2008. Lignin and polyphenols as allelochemicals. *Ind Crops Prod* 27: 144– 9.

9. Ignat I, Volf I, Popa VI. 2011. A critical review of methods for characterization of polyphenolic compounds in fruits and vegetables. *Food Chem* 126: 1821– 35.
10. Balasundram N, Sundram K, Samman S. 2006. Phenolic compounds in plants and agri-industrial by-products: antioxidant activity, occurrence, and potential uses. *Food Chem* 99: 191– 203.
11. Heim KE, Tagliaferro AR, Bobilya DJ. 2002. Flavonoid antioxidants: chemistry, metabolism, and structure-activity relationships. *J Nutr Biochem* 13: 572– 84.
12. Hollman PH, Katan MB. 1999. Dietary flavonoids: intake, health effects, and bioavailability. *Food Chem Toxicol* 37: 937– 42.
13. Robbins RJ. 2003. Phenolic acids in foods: an overview of analytical methodology. *J Agric Food Chem* 51: 2866– 87.
14. Bocco A, Cuvelier ME, Richard H, Berset C. 1998. Antioxidant activity and phenolic composition of citrus peel and seed extracts. *J Agric Food Chem* 46: 2123– 9.
15. Balasundram N, Sundram K, Samman S. 2006. Phenolic compounds in plants and agri-industrial by-products: antioxidant activity, occurrence, and potential uses. *Food Chem* 99: 191– 203.
16. Someya S, Yoshiki Y, Okubo K. 2002. Antioxidant compounds from bananas (*Musa cavendish*). *Food Chem* 79: 351– 4.
17. González-Centeno MR, Rosselló C, Simal S, Garau MC, López F, Femenia A. 2010. Physico-chemical properties of cell wall materials obtained from ten grape varieties and their byproducts: grape pomaces and stems. *LWT-Food Sci Technol* 43: 1580– 6.
18. González-Montelongo R, Lobo MG, González M. 2010. Antioxidant activity in banana peel extracts: testing extraction conditions and related bioactive compounds. *Food Chem* 119: 1030– 9.
19. Chang S, Tan C, Frankel EN, Barrett DM. 2000. Low-density lipoprotein antioxidant activity of phenolic compounds and polyphenol oxidase activity in selected clingstone peach cultivars. *J Agric Food Chem* 48: 47– 51.
20. Safdar MN, Kausar T, Jabbar S, Mumtaz A, Ahad K, Saddozai AA. 2016. Extraction and quantification of polyphenols from kinnow (*Citrus reticulata* L.) peel using ultrasound and maceration techniques. *J Food Drug Anal* 25(3): 488– 500.

21. Khoddami A, Wilkes MA, Roberts TH. 2013. Techniques for analysis of plant phenolic compounds. *Molecules* 18: 2328–75
22. Smith RM. 2003. Before the injection—modern methods of sample preparation for separation techniques. *J Chromatogr A* 1000: 3–27.
23. Hernández Y, Lobo MG, González M. 2009. Factors affecting sample extraction in the liquid chromatographic determination of organic acids in papaya and pineapple. *Food Chem* 114: 734–41.
24. Dorta E, Sogi DS. 2017. Value-added processing and utilization of pineapple by-products. In: MG Lobo, RE Paull, editors. *Handbook of pineapple technology, production, postharvest science, processing, and nutrition*. Oxford: John Wiley and Sons. p 196–220.
25. Dorta E, Lobo MG, González M. 2012a. Using drying treatments to stabilize mango peel and seed: effect on antioxidant activity. *LWT-Food Sci Technol* 45: 261–8
26. Zhou N., Chen H., Xi J., Yao D., Zhou Z., Tian Y., Lu X. Biochars with excellent Pb(II) adsorption property produced from fresh and dehydrated banana peels via hydrothermal carbonization. *Bioresour. Technol.* 2017;232:204–210.
27. Enniya I., Jourani A. Study of Methylene Blue Removal by a bio sorbent prepared with Apple peels. *J. Mater. Environ. Sci.* 2017;8:4573–4581.
28. Zaragoza M.D.L.L.Z., González-Reza R.M., Mendoza-Munoz N., Miranda-Linares V., Bernal-Couoh T.F., Mendoza-Elvira S., Quintanar-Guerrero D. Nanosystems in Edible Coatings: A Novel Strategy for Food Preservation. *Int. J. Mol. Sci.* 2018;19:705.
29. Wu Y., Luo Y., Wang Q. Antioxidant and antimicrobial properties of essential oils encapsulated in zein nanoparticles prepared by liquid-liquid dispersion method. *LWT.* 2012;48:283–290.
30. Prakash A., Baskaran R., Vadivel V. Citral nanoemulsion incorporated edible coating to extend the shelf life of fresh-cut pineapples. *LWT.* 2020;118:108851. Raghav P.K., Agarwal N., Saini M. Edible coating of fruits and vegetables: A review. *Int. J. Sci. Res. Mod. Edu.* 2016;1:188–204.
31. Ullah A., Abbasi N., Shafique M., Qureshi A.A. Influence of Edible Coatings on Biochemical Fruit Quality and Storage Life of Bell Pepper cv. “Yolo Wonder.” *J. Food Qual.* 2017;2017:1–11.

32. Alparslan Y., Metin C., Yapıcı H.H., Baygar T., Günlü A., Baygar T. Combined effect of orange peel essential oil and gelatin-coating on the quality and shelf life of shrimps. *J. Food Saf. Food Qual.* 2017;68:69–78.
33. Etxabide A., Urdanpilleta M., Gómez-Arriaran I., De La Caba K., Guerrero P. Effect of pH and lactose on cross-linking extension and structure of fish gelatin films. *React. Funct. Polym.* 2017;117:140–146.
34. Hanani Z.A.N., Yee F.C., Nor-Khaizura M. Effect of pomegranate (*Punica granatum L.*) peel powder on the antioxidant and antimicrobial properties of fish gelatin films as active packaging. *Food Hydrocoll.* 2019;89:253–259.
35. Shin S.H., Chang Y., Lacroix M., Han J. Control of microbial growth and lipid oxidation on a beef product using an apple peel-based edible coating treatment. *LWT.* 2017;84:183–188.
36. Al-Anbari I.H., Dakhel A.M., Adnan A. The effect of adding local orange peel powder to microbial inhibition and oxidative reaction within edible film component. *Plant Arc.* 2019;19:1006–1012.

Publisher's Note:

IRPG remains neutral with regard to jurisdictional claims in published maps and institutional affiliations. The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of IRPG and/or the editor(s). IRPG disclaims responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.